



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

A similar experiment with anhydrous succinic acid would produce succinate of baryta and a homologue of carbonic acid, the existence of which is also indicated by other considerations. It is premature to dwell upon this point; but in this direction also I have made some experiments.

II. "Notice of Researches on the Sulphocyanide and Cyanate of Naphtyl, conducted by VINCENT HALL, Esq." By A. W. HOFMANN, Ph.D., F.R.S. &c. Received August 10, 1858.


The transformation of phenylcarbamide and phenylsulphocarbamide under the influence of anhydrous phosphoric acid, respectively into cyanate and sulphocyanide of phenyl, an account of which I submitted to the Society several months ago, suggested the probability that the hitherto unknown cyanates and sulphocyanides of radicals similar to phenyl might be obtained by analogous processes.

To establish this point experimentally, Mr. Vincent Hall has examined, in my laboratory, the deportment of some of the derivatives of *naphthylamine* under the influence of agents capable of fixing ammonia and its analogues.


Mr. Hall has found that the crude naphthaline, such as it is obtained from the gas-works, submitted at once, without sublimation, to the action, first of fuming nitric acid, and subsequently of acetic acid and metallic iron, furnishes the naphthylamine sufficiently pure for these experiments. The crude product thus obtained was digested with bisulphide of carbon in order to convert it into *naphthylsulphocarbamide*.

By distilling naphthylsulphocarbamide with anhydrous phosphoric acid, Mr. Hall has obtained a beautiful crystalline compound of a faint but peculiar odour, readily fusible, easily soluble in alcohol and ether, insoluble in water.

The analysis of this compound has led to the formula




showing that it is in fact *sulphocyanide of naphthyl*, formed according to the equation:—




Boiled with an alcoholic solution of naphtylamine, this compound readily reproduces naphtylsulphocarbamide, which by its insolubility is easily distinguished and separated from the sulphocyanide.

Gently heated with phenylamine, the new sulphocyanide gives rise to the formation of a crystalline compound, of properties very similar to those of the naphtylsulphocarbamide. This new body is *phenyl-naphtyl-sulphocarbamide**, containing—




Naphtylcarbamide, as obtained by the action of potassa on the corresponding sulpho-compound, or by the distillation of oxalate of naphtylamine, is likewise powerfully attacked by anhydrous phosphoric acid. Among the products of distillation a compound is found, which, by its chemical properties, is readily identified as *cyanate of naphtyl*.




although the small quantity in which this body is produced—by far the greater amount of the naphtylcarbamide being charred by the action of anhydrous phosphoric acid—has hitherto prevented Mr. Hall from fixing the nature of the compound by an analysis.

* By the action of sulphocyanide of phenyl upon naphtylamine, I have obtained a crystalline compound very similar in its general characters to the body which Mr. Hall procures by the action of sulphocyanide of naphtyl on phenylamine. This substance likewise contains



for



Are these two bodies identical, or only isomeric? [A.W.H.]